(11) Publication number: 0 626 402 A3

(12)

## **EUROPEAN PATENT APPLICATION**

(21) Application number: 94303811.7

7.7

(51) Int. Cl.5: C08G 61/08

(22) Date of filing: 26.05.94

③ Priority: 27.05.93 US 68236 27.05.93 US 68240

15.12.93 US 167668

(43) Date of publication of application: 30.11.94 Bulletin 94/48

84 Designated Contracting States : BE DE ES FR GB IT NL

(8) Date of deferred publication of search report: 27.09.95 Bulletin 95/39

7) Applicant: AMOCO CORPORATION 200 East Randolph Drive Chicago Illinois 60601 (US)

Inventor: Nubel, Philip O.
 819 Tulip Lane
 Naperville, Illinois 60540 (US)

Inventor: Yokelson, Howard B. 1339 Brush Hill Circle Naperville, Illinois 60540 (US) Inventor: Moreland, Robert B. 29W225 Hartmand Drive Naperville, Illinois 60564 (US) Inventor: Bagheri, Vahid 6000 Oakwood Drive, 5H Lisle, Illinois 80532 (US) Inventor: Cohen, Steven A. 962 East Prarie Avenue Naperville, Illinois 60540 (US) Inventor: Bouslog, William G. 1517 West Jefferson Naperville, Illinois 60540 (US) Inventor: Behrends, Raymond T. 425 North Stewart Avenue Lombard, Illinois 60148 (US) Inventor: Nelson, James P. 1121 Kings Point Court Naperville, Illinois 60563 (US)

74 Representative: Schlich, George William et al Mathys & Squire 100 Grays Inn Road London WC1X 8AL (GB)

- Process for preparing linear monofunctional and telechelic difunctional polymers and compositions obtained thereby.
- This invention relates to a process for preparation of non-crosslinked linear monofunctional and telechelic difunctional unsaturated polymers wherein the functional groups are reactive terminal groups other than vinyl groups. The average functionality number of the monofunctional unsaturated polymers is at least 0.7, as determined by nuclear magnetic resonance spectroscopy (NMR). The average functionality number of the telechelic difunctional polymers is at least 1.6-1.7, as determined by NMR. Monofunctional olefins and difunctional olefins are reacted with cyclic olefins or unsaturated polymers to prepare difunctional polymers. The process is substantially free of side reactions comprising double bond migration and cyclization. The unsaturated functional polymers can be reacted in condensation, addition and transesterification reactions to prepare polyesgraft copolymers, polyurethanes, polyureas, polyamides, thermoplastic and therresins, block copolymers, exchange resins, adhesives, films, fibers, foams and flocculants.

P 0 626 402 A3



## EUROPEAN SEARCH REPORT

Application Number

	Citation of document with indicati	on, where anaronrists	Relevant	CLASSIFICATION OF THE
Category	of relevant passages		to claim	APPLICATION (Int.CL5)
A	EP-A-0 173 473 (BRITISH 5 March 1986 * page 12, line 27 - li			C08G61/08
A	EUROPEAN POLYMER JOURNA vol. 25, no. 7/8, 1989 pages 805-811, H.T.HO ET ALL. 'METATH OF SUBSTITUTED NORBORNE OF THE POLYMERS OF SOME * page 806; table 2 *	OXFORD GB, HESIS POLYMERIZATION NES:MICROSTRUCTURE	<del>-</del> .	
A .	MACROMOLECULES, vol. 20, 1987 EASTON Upages 778-782, P.A.PATTON ET ALL. 'FATTHE RATE OF TUNGSTEN HEXACHLORIDE/TETRAMETHY OLEFIN METHATHESIS POLY RING STRAIN IS NOT IMPO * page 1, paragraph 1 *	CTORS INFLUENCING  LTIN-CATALYSED  MERIZATION: MONOMER  RTANT'	i ja semante.	TECHNICAL FIELDS SEARCHED (Int. Cl.5)
,				C08G C08F
			÷	
			÷	
			•	
	The present search report has been dra	wn up for all claims		:
	Place of search	Date of completion of the search		Executant
	THE HAGUE	25 July 1995	Sti	enon, P
X : part	CATEGORY OF CITED DOCUMENTS  cularly relevant if taken alone cularly relevant if combined with another ment of the same category	I : theory or principle E : earlier patent doc after the filing da D : document cited for L : document cited for	ument, but publ te the application	ished on, or